

The unforgiving universe (semiclassicality in Loop Quantum Cosmology)

ILQG SEMINAR

15.02.2011

Tomasz Pawłowski

Department of Mathematics and Statistics

University of New Brunswick, Canada


work by:

W. Kamiński, TP: arXiv:1001.2663

A. Ashtekar, TP *unpublished*

The problem

- Loop quantization of the isotropic/homogeneous cosmological models \Rightarrow changes of the dynamics at near-Planck densities causing the Big Bounce.



Observation: States sharply peaked throughout the evolution.

- **Problem:** In numerical simulations one has to select an example of the state for evolution:

is the preservation of the semiclassicality robust?

- Addressing on the quantum level: Monte Carlo methods – probing the space of solutions via random samples (brute force approach).

Need to run large number of time-costly simulations!

- Any Alternative approach?

Alternatives

- If the system solvable analytically: One can find the relation between dispersions at distant future and past.
 - Affirmative answer in sLQC: *A. Corichi, P. Singh, arXiv:0710.4543*
States sufficiently sharply peaked initially remain so!
 - **Problem:** We need **exact solvability** of the system.
- Semiclassical dynamics: *M. Bojowald et al, arXiv:0911.4950, ...*
 - Choose canonical pair of variables X, P s.t. X, P and the momenta $G^{m,n} := \langle (\hat{X} - X)^m (\hat{P} - P)^n \rangle$ form closed algebra with evolution generator.
 - Capture the quantum dynamics as the EOMs for $G^{m,n}$.
 - **Problem:** Definiteness of the system of EOMs requires $|G^{m,n}| < \infty$: **states decay faster than polynomially for both X, P .**
If X, P chosen naively, for many systems such states may not exist!
- **The goal:** Flexible and reliable method of comparing the distant future and past states.
The means: The scattering picture.

Outline

An application to the simplest model:

- The flat FRW universe: specification.
- Geometrodynamical (WDW) vs LQC quantization.
- Method introduction for $\Lambda = 0$
 - WDW limit of LQC state.
 - The definition of the scattering picture.
 - An application: relation between dispersions.
- More complicated application: $\Lambda > 0$
 - WDW and LQC quantum system, deSitter limit.
 - The instantiations.
 - Appl: semiclassicality preservation between the cycles of evolution.
- Summary: Results and method properties.

An example model

Flat isotropic universe with massless scalar field.


- **Spacetime:** manifold $M \times \mathbb{R}$ where M is topologically \mathbb{R}^3 .
 $M \times \{t\}$ (where $t \in \mathbb{R}$) – homogeneous slices.
- **Metric:** $g = -dt^2 + a^2(t) {}^oq$
 - oq - flat fiducial metric ($dx^2 + dy^2 + dz^2$).
- **The treatment:**
 - The degrees of freedom (canonical variables):
 - Geometry: (v, b) , v - prop. to the volume of chosen fiducial region.
 - Matter: (ϕ, p_ϕ) , ϕ - scalar field value
 - **The quantization:**
 - **Matter:** standard Schrödinger representation
 - **Geometry:**
 - Wheeler-DeWitt: Schrödinger representation
 - **LQC:** methods of Loop Quantum Gravity
 - Nontrivial Hamiltonian constraint: Dirac program.
 - Scalar field used as an internal time

Wheeler-DeWitt quantization

All elements expressed in (c, p, ϕ, p_ϕ) . Standard quantization.

- **Kinematical Hilbert space:** $(v \propto \text{sgn}(p)|p|^{3/2})$
 - $\mathcal{H}^{\text{kin}} = \mathcal{H}^{\text{grav}} \otimes \mathcal{H}^\phi$, $\mathcal{H}^{\text{grav}} = L^2(\mathbb{R}, dv)$, $\mathcal{H}^\phi = L^2(\mathbb{R}, d\phi)$.
 - **Basic operators:** $(\hat{p}, \hat{c} \propto i\partial_p, \hat{\phi}, \hat{p}_\phi \propto i\partial_\phi)$.
 - **Basis: eigenstates** $(v|\hat{p} = v|, (\phi|\hat{\phi} = \phi|)$.
- **Quantum constraint:**
 $[\partial_\phi^2 \Psi](v, \phi) = -[\underline{\Theta}\Psi](v, \phi) := 12\pi G[(v\partial_v)^2 + v\partial_v + 1/4]\Psi(v, \phi)$.
- **Physical states:** $\Psi(v, \phi) = \int_{\mathbb{R}} dk \underline{\Psi}(k) \underline{e}_k(v) e^{i\omega\phi}$,
 $\underline{\Psi} \in L^2(\mathbb{R}, dk)$, $\omega = \sqrt{12\pi G|k|}$, $\underline{e}_k(v) = (1/\sqrt{2\pi v}) e^{ik \ln(v)}$.
- **Observables:**
 - $\hat{p}_\phi : \underline{\Psi}(k) \mapsto \hbar\omega(k)\underline{\Psi}(k)$,
 - $\ln(v)_{\phi_o} : \Psi(v, \phi) \mapsto e^{i\sqrt{\underline{\Theta}}(\phi-\phi_o)} \ln(v)\Psi(v, \phi_o)$.
- **Dynamics:**
 - **Two classes:** ever contracting and ever expanding.
 - The dispersions $\sigma_{\ln(v)_\phi}$ are **constant**.

WDW dynamics



LQC quantization

Geometry polymeric, matter standard, structure analogous

● Kinematics:

● Geometry space: $\mathcal{H}^{\text{grav}} = L^2(\mathbb{R}_{\text{Bohr}} d\mu_{\text{Bohr}})$.

● Geometry basis: $|v\rangle : \langle v|v'\rangle = \delta_{vv'}$.

● Basic operators: holonomies $h = e^{\int_{\gamma} A dx}$, fluxes $S = \int_S \star E d\sigma$.

● The constraint: reexpressed in terms of \hat{h} , \hat{S}

$$[\Theta\psi](v) = -f^+(v)\psi(v+4) + f^0(v)\psi(v) - f^-(v)\psi(v-4)$$

for large v : $f^{0,\pm}(v) \propto v^2$.

● The physical states: $\Psi(v, \phi) = \int_{\mathbb{R}^+} dk \tilde{\Psi}(k) e_k(v) e^{i\omega\phi}$
 $\tilde{\Psi} \in L^2(\mathbb{R}^+, dk)$, $\omega = \sqrt{12\pi G|k|}$, $\Theta e_k(v) = \omega^2(k) e_k(v)$.


● Dirac observables: analogous to WDW.

● The dynamics:

● in distant future and past **agreement with GR**,

● **bounce** in the Planck regime (energy densities).

LQC dynamics



WDW limit of LQC states

● $f^{o,\pm}$ - real: e_k are standing waves.

● **Observation:** converge to WDW standing waves.

$$e_k(v) = \underline{\psi}_k(v) + O(|\underline{e}_k(v)|(k/v)^2)$$
$$\underline{\psi}_k(v) := r(k)[e^{i\alpha(k)}\underline{e}_k(v) + e^{-i\alpha(k)}\underline{e}_{-k}(v)]$$

● **Comp. of LQC and WDW norms + self-adjointness of Θ :** $r(k) = 2$

● **Analytical proof:**

- reformulation of the diff. equation in the 1st order form,
- (local) decomp. of the LQC eigenf. in terms of WDW ones,
- asymptotic properties of the resulting transfer matrix.

● **Properties of the phase shifts:**

● **Analytic results for sLQC:** simple analytic form of the eigenfunctions in the momentum of v + stationary phase method

● $\alpha(k) = -k(\ln |k| - 1) - (3/4)\pi + o(k^0)$

● $\alpha'(k) = -\ln |k| + O(k^{-1} \ln |k|)$

● **Numerical results for: APS, sLQC, MMO:**

● $\alpha'(k) = -\ln |k| + O(k^{-2})$

● $|k\alpha''(k)| \leq 1$

The phase shifts


Properties analyzed analytically and numerically:

- **Analytic results for sLQC:** simple analytic form of the eigenfunctions in the momentum of v + stationary phase method
 - $\alpha(k) = -k(\ln |k| - 1) - (3/4)\pi + o(k^0)$
 - $\alpha'(k) = -\ln |k| + O(k^{-1} \ln |k|)$
- **Numerical results for: APS, sLQC, MMO:**
 - $\alpha'(k) = -\ln |k| + O(k^{-2})$
 - $|k\alpha''(k)| \leq 1$

Very regular behavior!


Phase shift properties

$$\alpha'(k).$$



Phase shift properties

$$k\alpha''(k).$$



The scattering picture

- WDW limit: $e_k(v) \rightarrow \underline{\psi}_k(v)$

$$\tilde{\Psi}(|k|) \mapsto \underline{\tilde{\Psi}}(k) = 2e^{i \operatorname{sgn}(k)\alpha(|k|)} \operatorname{sgn}(k) \tilde{\Psi}(|k|)$$

- Two components: $\underline{\tilde{\Psi}}_{\pm}(k) = \theta(\pm k) \underline{\tilde{\Psi}}(k)$

- The limits of observables:

$$\begin{aligned} \lim_{\phi \rightarrow \pm\infty} \langle \Psi | \ln(v)_{\phi} | \Psi \rangle &= \langle \underline{\Psi}_{\pm} | \ln(v)_{\phi} | \underline{\Psi}_{\pm} \rangle, \\ \lim_{\phi \rightarrow \pm\infty} \langle \Psi | \Delta \ln(v)_{\phi} | \Psi \rangle &= \langle \underline{\Psi}_{\pm} | \Delta \ln(v)_{\phi} | \underline{\Psi}_{\pm} \rangle =: \sigma_{\pm}. \end{aligned}$$

- Interpretation as a scattering process:

$$|\underline{\Psi}\rangle_{\text{in}} \mapsto |\underline{\Psi}\rangle_{\text{out}} = \hat{\rho} |\underline{\Psi}\rangle_{\text{in}}, \quad \underline{\tilde{\Psi}}_{\text{in}}(k) := \underline{\tilde{\Psi}}_{+}(k), \quad \underline{\tilde{\Psi}}_{\text{out}}(k) := \underline{\tilde{\Psi}}_{-}(k)$$


- The scattering matrix:

$$\rho(k, k') = (\underline{e}_k | \hat{\rho} | \underline{e}_{k'}) = e^{-i \operatorname{sgn}(k')\alpha(|k'|)} \delta(k + k').$$

- The transformation: total reflection

$$\underline{\tilde{\Psi}}(k) \mapsto U \underline{\tilde{\Psi}}(k) := e^{2i \operatorname{sgn}(k)\alpha(|k|)} \underline{\tilde{\Psi}}(-k)$$

The scattering



The dispersion growth

An application: comparizon of the dispersions of $\ln(v)_\phi$.

- Action on \mathcal{H}^{phy} : $\ln |\hat{v}|_\phi \tilde{\Psi} = [-i\partial_k - (\partial_k \omega(k)) \phi \hat{\mathbb{I}}] \tilde{\Psi}$

- The relation between action on limits: $\langle \hat{O} \rangle_\pm := \langle \underline{\Psi}_\pm | \hat{O} | \underline{\Psi}_\pm \rangle$

$$\langle -i\partial_k \rangle_- = \langle U^{-1}[-i\partial_k]U \rangle_+, \quad \langle \Delta[-i\partial_k] \rangle_- = \langle \Delta U^{-1}[-i\partial_k]U \rangle_+.$$

where $U^{-1}[-i\partial_k]U = -i\partial_k - 2\alpha'\mathbb{I}$.

- The effects on dispersions: ($\sigma_{A+B} \leq \sigma_A + \sigma_B$ - Schwartz ineq.)

$$\sigma_- \leq \sigma_+ + 2\langle \Delta\alpha'\mathbb{I} \rangle_+$$

- Estimate via dispersion in ω :

- General inequality:

$$\langle \Delta\alpha'\mathbb{I} \rangle_+^2 = \langle (\alpha'^2 - \langle \alpha' \rangle_+)^2 \mathbb{I} \rangle_+ \leq \langle (\alpha'^2 - \alpha'^*)^2 \mathbb{I} \rangle_+.$$

- The choice: $\alpha'^* = \alpha'(\exp(\lambda^*))$, $\lambda^* := \langle \ln(k) \rangle_+$ and props. of α'

give: $\langle \Delta\alpha'\mathbb{I} \rangle_+^2 \leq \langle (\ln(\hat{k}) - \lambda^*\mathbb{I})^2 \rangle_+ = \langle \Delta \ln(\hat{k}) \rangle_+^2 =: \sigma_\star^2$

- The final inequality:

$$\sigma_- \leq \sigma_+ + 2\sigma_\star$$

Summary

- **The results:** Devised method of comparing the properties of distant future and distant past states
 - does not require exact solvability
 - uses only asymptotic properties of phys. Hilbert space basis
 - is general (without restriction to particular types/shapes of states),
 - in genuinely quantum (no semiclassical approximations of any kind),
 - application to FRW with massless scalar:
general triangle inequalities on dispersions. Strict **upper bound** on eventual dispersion growth.

Just the upper bound, the actual dispersion may even shrink.

Universe's memory has to be indeed very sharp !

- generalization:
 - isotropic sector of Bianchi I
 - slightly weaker version: vacuum Bianchi I: arXiv:0906.3751
- further extension: $\Lambda > 0$ (see 2nd part).

$\Lambda > 0$ - *WDW model*

- Hamiltonian constraint:

$$-\partial_\phi^2 = \underline{\Theta}_o - \pi G \gamma^2 \Delta \Lambda \mathbb{I} = \underline{\Theta}_\Lambda$$

$\underline{\Theta}_\Lambda$ admits 1d family of selfadjoint extensions, labeled by $\beta \in U(1)$

- Each extension $\underline{\Theta}_{\Lambda\beta}$ has **continuous spectrum**: $\text{Sp}(|\underline{\Theta}_{\Lambda\beta}|) = \mathbb{R}^+$

- Physical states:

$$\begin{aligned} \underline{\Psi}(v, \phi) &= \int_0^\infty dk \tilde{\Psi}(k) e_k^\beta(v) e^{i\omega\phi}, \quad \omega = \sqrt{12\pi G} k \\ e_k^\beta(v) &= \frac{1}{\sqrt{|v|}} [c_1(\beta, \Lambda, k) H_{ik}^{(1)}(av) + c_2(\beta, \Lambda, k) H_{ik}^{(2)}(av)], \end{aligned}$$

where $a = \sqrt{\frac{\gamma^2 \Delta \Lambda}{12\pi G}}$ and $H^{(1)}, H^{(2)}$ - Hankel functions.

- For each extension all e_k^β have **common** asymptotics

$$e_k^\beta = N(\Lambda, \beta, k) |v|^{-1} \cos(\Omega(\Lambda) |v| + \sigma(\Lambda, \beta)) + O(|v|^{-3/2})$$

- **Dynamics**: follows analytically extended classical trajectory.

WDW: observables

- Observables: analogous to $\Lambda = 0$ case.

Problem: the analog of $|v\rangle_\phi$ leads **outside of** \mathcal{H}^{phy} !

- **Failure of semiclassical treatment:**

- If one selects canonical pair v, b the requirement $\langle (|\hat{v}\rangle_\phi - \langle v|\phi\rangle)^n \rangle < \infty$ on some open $\phi \in \mathcal{O}$ implies

$$\int dk \tilde{\Psi}(k) N(\Lambda, \beta, k) e^{i\omega\phi} = 0, \quad \forall \phi \in \mathcal{O}$$

- Since $N(\Lambda, \beta, k) \propto \sqrt{k}$ we have: $\tilde{\Psi}(k) \sim k^{-3/2} \Rightarrow \langle \Delta p_\phi \rangle = \infty$.

- For $|v| \ll 1$ $e_k^\beta \approx e^{ik \ln |v|}$, thus at early times


$$\langle \Delta p_\phi \rangle = \infty \Rightarrow \langle \Delta b|_\phi \rangle = \infty$$

Impossible to built states well behaving in both v and b even for a short time!


- **Solution:**

- Compactify v , for example use $\theta_a = \arctan(|v|/a)$ or
- Use truly measurable quantities, like Hubble parameter H or energy density ρ .

WDW dynamics for $\Lambda > 0$



WDW dynamics for $\Lambda > 0$



$\Lambda > 0$: *LQC model*

- Allowed value $\Lambda \in [0, \Lambda_c]$, where $\Lambda_c = 8\pi G\rho_c$

- The constraint (sLQC prescription):

$$-\partial_\phi^2 = \Theta_o - \pi G\gamma^2 \Delta\Lambda\mathbb{I} = \Theta_\Lambda$$

Θ_Λ admits 1d family of selfadjoint extensions, labeled by $\beta \in U(1)$

- Each extension $\Theta_{\Lambda\beta}$ has **discrete spectrum**: $\text{Sp}(\Theta_{\Lambda\beta}) = \{\omega_n^2\}$

where $\omega_n = [\sqrt{12\pi^3 G}/f_1(\Lambda)] \cdot k_n$,

$$\tan(g(\Lambda)k_n) + \tanh[(\pi - g(\Lambda))k_n] \tan(\beta) = 0, \quad g(\Lambda) \in [0, \pi]$$

$$\Rightarrow \omega_n = (n\pi - \beta)/f_2(\Lambda) + O(e^{-2\pi n(\pi - g(\Lambda))/g(\Lambda)})$$

- Physical states:**

$$\Psi(v, \phi) = \sum_{n=0}^{\infty} \tilde{\Psi}_n e_n^\beta(v) e^{i\omega_n \phi}$$


- For each extensions common leading order asymptotics

$$e_n^\beta = N_n(\Lambda, \beta) |v|^{-1} \cos(\Omega(\Lambda)|v| + \sigma(\Lambda, \beta)) + O(|v|^{-3/2})$$


- Dynamics:**

- Agreement with GR for low energies.
- Bounce in Planck regime.

LQC dynamics for $\Lambda > 0$



LQC dynamics for $\Lambda > 0$



2nd order asymptotics

- Analysis of the transfer matrix asymptotics:

$$e_n^\beta(v) = N_n [e^{i\alpha} e_n^+(v) + e^{-i\alpha} e_n^-(v)] + O(v^{-3}) \quad (\star)$$

$$e_n^\pm = |v|^{-1} e^{\pm i\Omega|v|} \cdot e^{\pm i\kappa(n, \Lambda, \beta)/|v|}, \quad 1/|v| \approx (1/a)(\theta_a - \pi/2)$$

where: $\cos(4\Omega(\Lambda)) = 1 - 2\Lambda/\Lambda_c =: 1 - 2\lambda$,

$$\kappa(n, \Lambda, \beta) = \frac{3\pi G(1-2\lambda) + \omega_n^2}{12\pi G\sqrt{\lambda(1-\lambda)}} =: A\omega_n^2 + B$$

- Limit spaces:

- “Standing wave form” of e_n^β :
split onto incoming and outgoing components.
- For each comp. $e^{\pm i\Omega|v|}$ is a global rotation, only e_n^\pm relevant.
- e_n^\pm form wave packet regular in θ_a . *Schrödinger type rather than Klein-Gordon.*

- Transformation into limit spaces:

- Define spaces \mathcal{H}^\pm spanned by e_n^\pm with IP inherited from \mathcal{H}^{phy} through the limit (\star) .
- $\mathcal{H}^{\text{phy}} \ni \tilde{\Psi}_n \mapsto \Phi_n^\pm := \tilde{\Psi}_n M_n \in \mathcal{H}^\pm$

WDW limit states

- Taking the limit of the mass gap $\Delta \rightarrow 0$ one can derive analogous 2nd order limit in WDW theory.
 - As $\Omega(\Lambda)$ depends implicitly on Δ to ensure uniform convergence one has to allow for flow $\Lambda = \Lambda(\Delta)$.
 - **Result:** LQC wave packet has the WDW limit spanned by \underline{e}_k such that:

$$\underline{e}_k^\beta(v) = N(k)[e^{i\alpha}\underline{e}_k^+(v) + e^{-i\alpha}\underline{e}_k^-(v)] + O(v^{-3})$$

where

$$\underline{e}_k^\pm = |v|^{-1} e^{\pm i\Omega|v|} \cdot e^{\pm i\kappa(k,\Lambda,\beta)/|v|}$$

with the same Ω and $\kappa = \kappa(\omega = \sqrt{12\pi Gk})$

however the WDW model corresponds to the cosmological constant

$$\underline{\Lambda}/\Lambda_c = \underline{\lambda} = \arccos(1 - 2\lambda).$$

- **Construction of the limit spaces $\underline{\mathcal{H}}^\pm$** analogous to LQC.

Again Schrödinger wave packets near $\theta = \pi/2$.

- If the wave packet sharply peaked about $\theta = \pi/2$ then (up to higher order corrections)

$$\langle \Delta | \hat{\theta}_a | \phi \rangle_{\text{WDW}} = \langle \Delta | \hat{\theta}_a | \phi \rangle_{\text{lim}}$$

Instantiations


● The continuous limit:

- We identified the relations between \mathcal{H}^{phy} , $\underline{\mathcal{H}}^{\text{phy}}$ and the appropriate limit spaces.
- To build $\mathcal{H}^{\text{phy}} \leftrightarrow \underline{\mathcal{H}}^{\text{phy}}$ we need $\mathcal{H}^{\pm} \leftrightarrow \underline{\mathcal{H}}^{\pm}$.
- **Problem:** Identification of Φ_n and $\Phi(k)$ will produce zero norm WDW states!

● The instantiations:

- Fix the moment $\phi = \phi_o$. By rotation $\tilde{\Psi}(k) \mapsto \tilde{\Psi}(k)e^{i\omega\phi_o}$ one can bring it to $\phi = 0$.
- On $\underline{\mathcal{H}}^{\pm}$ the operator $\hat{x} := \hat{\theta}_a - \pi/2$ takes the form $\hat{x} = \frac{ia}{2A\omega} \partial_\omega$
- On \mathcal{H}^{\pm} one can build “ $\sin(cx)/c$ ” oper. via $h : [h\tilde{\Phi}]_n = \tilde{\Phi}_{n-1}$
- $$\sin(cx)/c = \frac{ai}{2A[\Delta\omega](\omega + [\Delta\omega]/2)} [h - h^{-1}] + O(e^{-a\omega}),$$
 where $\Delta\omega = \lim_{n \rightarrow \infty} [\omega_n - \omega_{n-1}]$.
- If at ϕ_o state sharply peaked about $x = 0$: $\sin(cx)/c$ - good replacement of \hat{x}
- **Instantiation:** Interpolation of $\tilde{\Phi}_n$ s.th. actions of \hat{x}^n and $[\sin(cx)/c]^n$ agree up to $n = 2$.

Instantiations + scattering



The scattering, decoherence

The picture:

- Instantiations provide identification of quasi-periodic state of LQC with non-periodic WDW one at given time ϕ_o .
- Once can build a sequence of instantiations at ϕ_n where $\langle \theta \rangle \approx \pi/2$.
- Transfer $\phi_n \rightarrow \phi_{n+1}$: scattering of WDW state into another.
- Evolution: sequence of scatterings (determined by instantiations).
- Since $\omega_n = (n\pi - \beta)/f_2(\Lambda) + \delta\omega_n$ one cycle corresponds to

$$\tilde{\Psi}_n \mapsto e^{2\pi i \delta\omega_n} \tilde{\Psi}_n =: U \tilde{\Psi}_n$$

Application:

- $\langle \Delta \sin(cx)/c \rangle_{\mathcal{H}^{\text{phy}}} = \langle \Delta \sin(cx)/c \rangle_{\mathcal{H}^{\pm}} + O(v^{-3})$.
- After $N \gg 1$ cycles

$$\langle \Delta \sin(cx)/c \rangle_{\phi_o + N\Delta\phi} \leq \langle \Delta \sin(cx)/c \rangle_{\phi_o} + N \langle \Delta \frac{2\pi \partial_\omega \delta\omega}{\omega} \rangle$$

where the last term $\langle \Delta \frac{2\pi \partial_\omega \delta\omega}{\omega} \rangle \leq C \langle \Delta \frac{2\pi e^{-a\omega}}{\omega} \rangle$.

Summary

- Scattering picture **successfully extended** to the model with $\Lambda > 0$.
- Evolution between pure deSitter epochs - chain of Wheeler-DeWitt universes consecutively scattered one into another.
- The instantiation procedure allowed to relate the dispersion in compactified volume θ_a of the LQC state and its WDW limit at given moment ϕ_o (in pure deSitter epoch).
- The scattering corresponds to unitary rotation by $e^{2\pi i \delta\omega_n}$, where the deviation $\delta\omega_n$ decays exponentially.
- **Consequence:** The decoherence of the state between large number N of pure deSitter epochs is **bounded from above** by (up to a known constant) the dispersion of the operator $N e^{-a\omega} / \omega$ infinitesimal for the states peaked about large p_ϕ^* .

Appendix: The transfer matrix method

- $f^{o,\pm}$ - real: e_k are standing waves.
- **Observation:** converge to WDW standing waves.
- **Verification:** transfer matrices

- 1st order form of the difference equation:

$$\vec{e}_k(v) = \begin{bmatrix} e_k(v) \\ e_k(v-4) \end{bmatrix}, \quad A(v) = \begin{bmatrix} \frac{f_o(v) - \omega^2(k)}{f_+(v)} & -\frac{f_-(v)}{f_+(v)} \\ 1 & 0 \end{bmatrix}$$

$$\vec{e}_k(v+4) = A(v)\vec{e}_k(v)$$

- Expressing in WDW basis:

$$\vec{e}_k(v) = B(v)\vec{\chi}_k(v), \quad B(v) := \begin{bmatrix} \underline{e}_k(v+4) & \underline{e}_{-k}(v+4) \\ \underline{e}_k(v) & \underline{e}_{-k}(v) \end{bmatrix}.$$

- Final form: $\vec{\chi}_k(v+4) = B^{-1}(v)A(v)B(v-4)\vec{\chi}_k(v) =: M(v)\vec{\chi}_k(v)$

- **Limit of the transfer matrix:** $M(v) = \mathbb{I} + O(v^{-3}) \Rightarrow$

$$e_k(v) = \underline{\psi}_k(v) + O(|\underline{e}_k(v)|(k/v)^2)$$

$$\underline{\psi}_k(v) := r(k)[e^{i\alpha(k)}\underline{e}_k(v) + e^{-i\alpha(k)}\underline{e}_{-k}(v)]$$

- **Comparizon of norms:** $r(k) = 2$

Appendix: Comparizon between the norms

● Evolution: mapping $\mathbb{R} \ni \phi \mapsto \psi(\cdot) := \Psi(\cdot, \phi) \in \mathcal{H}^{\text{grav}}$

● Inner products:

$$\langle \psi | \chi \rangle = \sum_{\mathcal{L}_0^+} \overline{\psi(v)} \chi(v), \quad \langle \underline{\psi} | \underline{\chi} \rangle = \int_{\mathbb{R}^+} dv \overline{\underline{\psi}(v)} \underline{\chi}(v)$$

● Distributional estimates:

● splitting the domain:

$$X(k, k') := \langle e_{k'} | e_k \rangle = \sum_{\mathcal{L}_0^+ \cap [1, \infty]} \overline{e_{k'}(v)} e_k(v)$$

● extracting WDW limits:

$$X(k, k') = + \sum_{\mathcal{L}_0^+ \cap [1, \infty]} [\overline{\underline{\psi}_{k'}(v)} \underline{\psi}_k(v) + O(v^{-5/2})]$$

● estimating the sum by the integral:

$$\sum \overline{\underline{\psi}_{k'}(v)} \underline{\psi}_k(v) = (1/4) \int_{[1, \infty[} dv \left[\overline{\underline{\psi}_{k'}(v)} \underline{\psi}_k(v) + O(v^{-3/2}) \right]$$

● relation: $\int_{\mathbb{R}^+} dx e^{ikx} = \frac{1}{2} \left(\int_{\mathbb{R}} dx e^{ikx} - \frac{i}{\pi k} \right)$

● Final relation: $X(k, k') = (r^2(k)/8) \langle \underline{\psi}_k | \underline{\psi}_{k'} \rangle + F(k, k')$

● Orthonormality: $F(k, k') = 0, r(k) = 2.$

Appendix: The limits of observables

Argumentation for past limit, problem symmetric in time.

- Assumed localization in $\ln |v|_\phi$ of WDW limit: $\langle \ln |v|_\phi \rangle < \infty$,
 $\langle \Delta \ln |v|_\phi \rangle < \infty$.
- Past wave packet follows the trajectory $\bar{x}(\phi) = x_o - \beta(\phi - \phi_o)$,
where $\beta = \sqrt{12\pi G}$.
- $f(v)_\phi$ - multiplication operator on ID's $\Psi(\cdot, \phi) \in \mathcal{H}^{\text{grav}}$,
expectations - local sums: $\langle f(v)_\phi \rangle = \sum_{\mathcal{L}_0^+} f(v) |\Psi(v, \phi)|^2$,
analogous situation in WDW.
- introduce $\tilde{x}(\phi) = x_o - (\beta/2)(\phi - \phi_o)$ and split the local sums along it.
- the following properties:
 - falloff conditions due to localization: parts for $x > \tilde{x}$ converge
(sufficiently fast) to complete sums.
 - relation between LQC and WDW norms: convergence of WDW
and LQC partial sums for states localized in k .

imply that:

$$\lim_{\phi \rightarrow -\infty} \langle \Psi | \ln(v)_\phi | \Psi \rangle = \langle \underline{\Psi}_- | \ln(v)_\phi | \underline{\Psi}_- \rangle,$$
$$\lim_{\phi \rightarrow -\infty} \langle \Psi | \Delta \ln(v)_\phi | \Psi \rangle = \langle \underline{\Psi}_- | \Delta \ln(v)_\phi | \underline{\Psi}_- \rangle.$$